arXiv:cond-mat/0606705v1 [cond-mat.str-el] 28 Jun 2006

Impurity Entanglement Entropy and the Kondo Screening Cloud

Erik S. Sørensen,[1] Ming-Shyang Chang,[2] Nicolas Laflorencie,[2] and Ian Affleck[2]
[1]*Department of Physics and Astronomy, McMaster University, Hamilton, ON, L8S 4M1 Canada*
[2]*Department of Physics & Astronomy, University of British Columbia, Vancouver, B.C., Canada, V6T 1Z1*
(Dated: February 6, 2008)

The screening of an impurity spin by conduction electrons is associated with the formation of a large Kondo screening cloud, of size ξ_K. We study the quantum entanglement between a region of size r surrounding the impurity and the rest of the sample, (of total size $\mathcal{R}$) using Density Matrix Renormalization Group and analytic methods. The corresponding "impurity entanglement entropy", S_{imp}, is shown to be a universal scaling function of r/ξ_K and $r/\mathcal{R}$. We calculate this universal function using Fermi liquid theory in the regime $\xi_K \ll r$.

PACS numbers: 03.67.Mn,75.30.Hx,75.10.Pq

Quantum entanglement of many body wave-functions is receiving much attention of late due to its quantum information (QI) theory applications [1], its importance to computational algorithms [2] and the new perspective that it offers on critical behavior of condensed matter systems [3, 4]. Very recently, entanglement has been investigated in some quantum impurity models [5]. The Kondo Hamiltonian [6], a famous quantum impurity model, was originally proposed to represent a single impurity spin interacting with conduction electrons in a metal. The Hamiltonian is:

$$H = \int d^3r[\psi^\dagger(-\nabla^2/2m)\psi + J_K\delta^3(r)\psi^\dagger(\vec{\sigma}/2)\psi\cdot\vec{S}], \quad (1)$$

where $\psi(\vec{r})$ is the electron annihilation operator (with spin-index suppressed), the S^a are $S = 1/2$ spin operators and the Kondo coupling, $J_K > 0$. Due to the δ-function in Eq. (1), we may expand $\psi(\vec{r})$ in spherical harmonics and only the s-wave couples to the impurity, yielding an effective one dimensional ($1D$) model. From spin-charge separation in one dimension, it can be seen that the universal low energy long distance physics of the Kondo model also occurs in the $J_1 - J_2$ family of gapless antiferromagnetic $S = 1/2$ Heisenberg spin chain models for $J_2 \leq J_2^c \simeq 0.2412$, with the coupling of the end spin to its neighbors, J'_K, weaker than the other couplings [7]:

$$\begin{aligned} H = \; & J'_K\left(\vec{S}_1\cdot\vec{S}_2 + J_2\vec{S}_1\cdot\vec{S}_3\right) + \\ & \sum_{j=2}^{\mathcal{R}-1}\vec{S}_j\cdot\vec{S}_{j+1} + J_2\sum_{j=2}^{\mathcal{R}-2}\vec{S}_j\cdot\vec{S}_{j+2}. \end{aligned} \quad (2)$$

The Kondo coupling, J_K (or J'_K), is found to grow large at low energies under renormalization and the ground state is a highly non-trivial many body state in which the impurity spin is sometimes heuristically thought of as being screened by a conduction electron in a large orbital of size

$$\xi_K = v/T_K. \quad (3)$$

Here v is the velocity of the low energy excitations (fermions or spin-waves) and T_K is the energy scale at which the renormalized Kondo coupling becomes large. For weak bare coupling, $\xi_K \propto e^{\alpha/J_K}$ where α is a constant. Many physical observables were shown to vary on the characteristic length scale, ξ_K [8, 9, 10].



FIG. 1: (a) Schematic picture of the two regions A,B around a Kondo Impurity. (b) $1D$ analogue of the Kondo problem, with $\mathcal{R}$ interacting qubits along an antiferromagnetic chain. (c) Total entanglement entropy, $S(J'_K, r, \mathcal{R})$ for a 102 site spin chain at J_2^c, with a $J'_K = 0.3$ Kondo impurity along with $S(J'_K = 1, r-1, \mathcal{R}-1)$ (no impurity). Uniform parts (solid lines) and the resulting S_{imp}.

The purpose of this paper is to investigate the interplay of the Kondo screening cloud and quantum entanglement occurring in the models Eqs. (1-2). In a QI context we could think of the three dimensional ($3D$) fermionic model, Eq. (1), as representing a single qubit coupled to a dissipative environment. Alternatively, the $1D$ spin model, Eq. (2), corresponds to a chain of qubits with the end qubit more weakly coupled than the rest. $S = 1/2$ chains are receiving considerable attention from the QI community in the context of perfect state transfer [11]. In the 3D fermionic model, we imagine separating space into a spherical region, A, of radius r surrounding the impurity spin and the region B, exterior to the sphere as in Fig. 1(a). Tracing out the region B from the (pure) ground state density matrix, we obtain the reduced density matrix $\rho(r)$ for A. The entanglement entropy is defined as its Neumann entropy $S = -\mathrm{tr}[\rho(r)\ln\rho(r)]$ [12]. In the $1D$ spin chain version, we consider the entanglement of the first r spins with the remainder of the chain as in Fig. 1(b). For the $1D$ spin chain at J_2^c we show the resulting entanglement entropy in Fig. 1(c) for $J'_K = .3$ and with no impurity for $J'_K = 1$. Note that, in both cases, there is a strong even-odd alternation of S with r. This effect was studied in [13] for the case of uniform coupling, $J'_K = 1$. In the present paper we focus on the uniform part, which can be defined at $r \gg 1$ by:

$$S(r, \mathcal{R}) = S_U(J'_K, r, \mathcal{R}) + (-1)^r S_A(J'_K, r, \mathcal{R}), \qquad (4)$$

where $S_U(J'_K, r, \mathcal{R})$ and $S_A(J'_K, r, \mathcal{R})$ are slowly varying functions or r. There is also a strong alternation with the parity of $\mathcal{R}$ and we study both $\mathcal{R}$ even and odd. The uniform part consists of a bulk part, $S_{U0}(r, \mathcal{R}) \approx (1/6)\ln[\mathcal{R}\sin(\pi r/\mathcal{R})]$ + constant, independent of J'_K [4, 13, 14] and an impurity term that we define precisely as:

$$S_{imp}(J'_K, r, \mathcal{R}) \equiv S_U(J'_K, r, \mathcal{R}) - S_U(1, r-1, \mathcal{R}-1). \qquad (5)$$

Here we have subtracted S_U when the impurity is absent. See Fig. 1(c). There is also an impurity term in S_A but we do not address this here. We focus here on calculating S_{imp} for the spin chain model but the same universal scaling function also describes the additional entanglement entropy due to the impurity in the 3D Kondo model [15]. We find that $S_{imp}(J'_K, r, \mathcal{R})$ appears to be a scaling function, depending only on the ratios of characteristic lengths when r, $\xi_K \gg 1$:

$$S_{imp}(J'_K, r, \mathcal{R}) = S_{imp}(r/\xi_K, r/\mathcal{R}). \qquad (6)$$

Note that there are actually two different scaling functions for the cases of $\mathcal{R}$ even or odd. We have verified this scaling form over a wide range of parameters using the Density Matrix Renormalization Group (DMRG) keeping from 256 to 1024 states (Fig. 2). In the limit $\xi_K \ll r$ we have calculated the universal scaling function explicitly using conformal field theory (CFT) methods based on Nozières' local Fermi liquid theory (FLT) [16, 17].

Numerical Results. We have determined S_{imp} for a range of values of J'_K, r and $\mathcal{R}$ using DMRG on the spin chain model, Eq. (2). In order to avoid well-known logarithmic corrections due to a marginally irrelevant (bulk) operator [19] we tune J_2 to the critical point $J_2^c \approx .2412$, where the marginal coupling constant vanishes [7]. We then extract S_U from the DMRG data with high precision using a 7-point formula. In Fig. 2 we show a scaling plot of S_{imp} for several fixed $r/\mathcal{R}$ allowing for a determination of $\xi_K(J'_K)$ and validating the scaling form, Eq. (6).



FIG. 2: Universal scaling plot of S_{imp} for fixed $r/\mathcal{R}$, (a) for $\mathcal{R}$ *even*, (b) for $\mathcal{R}$ *odd*. DMRG results for the $J_1 - J_2$ chain at J_2^c for various couplings J'_K. The lines marked $\pi\xi_K/(12r)$ are the FLT prediction: Eq. (10). (c): the location of the maximum, $(r/\xi_K)_{\max}$, of S_{imp} for odd $\mathcal{R}$, plotted versus $r/\mathcal{R}$.

For odd $\mathcal{R}$, $S(0, r, \mathcal{R}) - S(1, r-1, \mathcal{R}-1)$ is exactly zero, since the decoupled impurity at $j = 1$ has no entanglement with the rest of the chain in the $S = 1/2$ ground state in which it is fully polarized. This implies that $S_{imp}(0, r, \mathcal{R}) = 0$ for $\mathcal{R}$ odd. On the other hand, for $\mathcal{R}$ even, $S_{imp}(0, r, \mathcal{R})$ is not zero, as shown in Fig. 3. Thus the presence of the impurity spin, even when there is no term in the Hamiltonian coupling it to the rest of the system, alters S. The reason this is possible is because, with $\mathcal{R}$ even and $J'_K = 0$, the ground state is 4-fold degenerate, corresponding to the impurity spin and the rest of the chain independently having $S^z = \pm 1/2$. The ground state that we consider in this case is the spin singlet state. This is necessary to connect to the limit $J'_K \to 0^+$. $S(0, r, \mathcal{R})$ is then clearly different from $S(1, r-1, \mathcal{R}-1)$ with a resulting non-zero $S_{imp}(0, r, \mathcal{R})$ for $\mathcal{R}$ even [15]. We expect that, in the limit $r/\mathcal{R} \to 0$, $S_{imp}(r/\xi_K)$ becomes the same for even and odd R, since adding one more site to an infinite chain should not have any effect on the entanglement entropy. Our data suggests that $\lim_{r/\xi_K \to 0} \lim_{r/R \to 0} S_{imp}(r/\xi_K, r/\mathcal{R}) = \ln 2$, for either parity of $\mathcal{R}$. We note that the presence of a term in the entanglement entropy at a conformally invariant boundary fixed point, corresponding to the zero temperature thermodynamic impurity entropy, $\ln g$ was pointed out in Ref. [4]. For the single channel Kondo model $\ln g = \ln 2$ at the weak coupling fixed point where the impurity is unscreened, and $\ln g = 0$ at the strong coupling fixed point where it is screened. The thermodynamic impurity entropy decreases monotonically from $\ln 2$ to 0 with decreasing T/T_K. The impurity entanglement entropy behaves the same way with increasing r/ξ_K.



FIG. 3: DMRG results using spin-inversion for $S_{\rm imp}(0,r,\mathcal{R})$ for $\mathcal{R}$ even. Inset: DMRG results for $S_{\rm imp}(1,r,\mathcal{R})$ for $\mathcal{R}$ odd and even. Data for $\mathcal{R} = 100, 101$ and $\mathcal{R} = 400, 401$ are indistinguishable. $S_{\rm imp}(1,r,\mathcal{R})$ appears to vanish in the scaling limit $r \to \infty$ with $r/\mathcal{R}$ held fixed.

A heuristic picture giving the general features of $S_{imp}(r/\xi_K, r/\mathcal{R})$, for both even and odd $\mathcal{R}$, can be derived from a resonating valence bond view of the ground state [18]. We loosely identify S_{imp} with $\ln 2\times$ the probability of an "impurity valence bond" (IVB) stretching from the impurity site into region B. (See Fig. 4.) For $J'_K = 0$ and even $\mathcal{R}$, all lengths for the IVB are equally probable so the naive picture predicts that S_{imp} decreases linearly from $\ln 2$ to 0 with increasing r/R, in rough agreement with the DMRG data in Fig. 3. We expect the typical length of the IVB to be ξ_K (when $\xi_K < \mathcal{R}$), leading to the monotonic decrease of S_{imp} with increasing r/ξ_K, for $\mathcal{R}$ even, and its vanishing as $r/\xi_K \to \infty$. For $\mathcal{R}$ odd, the ground state contains one unpaired spin. When $\xi_K = \infty$ this unpaired spin is the impurity, implying no IVB and hence $S_{imp} = 0$. As ξ_K decreases the probability of having an IVB increases, since the unpaired spin becomes more likely to be at another site, due to Kondo screening, but the average length of the IVB, when it is present, decreases. These two effects trade off to give S_{imp} a maximum when $\xi_K \approx R$, in good agreement with Fig. 2 inset, at which point the probability of having an IVB becomes $O(1)$ but the decreasing average size of the IVB starts to significantly decrease the probability of it stretching into region B.



FIG. 4: Typical valence bond configurations for $\mathcal{R}$ even (a) and $\mathcal{R}$ odd (b) where the unpaired spin is on the 7th site.

FLT calculation. The non-interacting part of the 1D (s-wave) fermion Hamiltonian can be written as the integral over r of spin and charge energy densities for left and right-movers, $\mathcal{H}_{s,L/R}$ and $\mathcal{H}_{c,L/R}$. At energy scales $\ll T_K$ the impurity spin is screened and does not appear in the low energy effective Hamiltonian. Irrelevant local interactions, at $r = 0$, not involving the impurity spin appear instead [17]. The most relevant of these can be written:

$$H_{int} = -(\xi_K \pi)\mathcal{H}_{s,L}(0). \tag{7}$$

Various long distance, low energy properties can be calculated in an expansion in ξ_K/r and $E\xi_K/v$ at distances $r \gg \xi_K$ and energy scales $E \ll v/\xi_K$. For instance, the zero temperature impurity susceptibility is $\chi(T = 0) = \xi_K/(4v)$. In the limit $\xi_K \ll r$, we can calculate S_{imp} using the FLT interaction of Eq. (7) in lowest order perturbation theory in ξ_K. This calculation is easily done by CFT methods. We first express trρ^n in terms of the partition function, $Z_n(r)$ on an n-sheeted Riemann surface, $\mathcal{R}_n$, with the sheets joined at the cut extending from r to $\mathcal{R}$ [4, 14]. The entanglement entropy is obtained by means of the replica trick:

$$S = -\lim_{n\to 1}\frac{d}{dn}[Z_n/Z^n]. \tag{8}$$

(Z is the partition function on the ordinary complex plane.) The correction to Z_n of first order in ξ_K is:

$$-\delta Z_n = -(\xi_K \pi) n \int_{-\infty}^{\infty} d\tau \langle \mathcal{H}_{s,L}(\tau, 0)\rangle_{\mathcal{R}_n}. \tag{9}$$

$\mathcal{H}_{s,L} = T/(2\pi)$ where $T(\tau, x)$ is the conventionally normalized energy-momentum tensor for the $c = 1$, free boson conformal field theory corresponding to the spin excitations of the original free fermion model. It obeys [4]: $\langle T(\tau,0)\rangle_{\mathcal{R}_n} = v\Delta_n(2ir)^2/[(v\tau+ir)^2(v\tau-ir)^2]$, with $\Delta_n = (c/24)(1-n^{-2})$. Both the integral in Eq. (9) and the limit in eq. (8) are trivial, yielding (see Fig 2):

$$S_{imp} = \pi\xi_K/(12r). \tag{10}$$

Our FLT calculation of S_{imp} is easily extended to finite $\mathcal{R}$. In this case, $\langle T\rangle$ is given as above with the substitution: $v\tau \pm ir \rightarrow (2\mathcal{R}/\pi)\sinh[\pi(v\tau \pm ir)/(2\mathcal{R})]$, and $2r \rightarrow (2\mathcal{R}/\pi)\sin[\pi r/(\mathcal{R})]$. The integral in Eq. (9) can again be done exactly, yielding:

$$S_{imp} = [\pi\xi_K/(12\mathcal{R})][1 + \pi(1 - r/\mathcal{R})\cot(\pi r/\mathcal{R})]. \quad (11)$$

We note that this result can be simply obtained from $S_{U0} = (1/6)\ln[\mathcal{R}\sin\pi r/\mathcal{R})]$ by shifting both r and R by $\pi\xi_K/2$ and Taylor expanding to first order in the small quantities ξ_K/r and ξ_K/R. Excellent agreement between this result and DMRG data is shown in Fig. 5(a) allowing for a determination of $\xi_K(J'_K)$.

In Figs. 2,5(a) we have determined $\xi_K(J'_K)$ by requiring good data collapse. It is an important consistency check on our scaling hypothesis to also determine $\xi_K(J'_J)$ independently by more conventional methods. Using the beta-function to cubic order we find that [15]: $\xi_K \propto \exp(1.376/J'_K)/\sqrt{J'_K}$. [To derive the 1.376 factor we needed the spin-wave velocity, $v \approx 1.1699$ and the coefficient of proportionality between $\vec{S}_2 + J_2^c\vec{S}_3$ and the free fermion spin density at the impurity location, which was determined by comparing the end to end spin correlation function (decaying as $1/\mathcal{R}^2$) in the spin chain and $1D$ free fermion models.] This formula is only valid at very weak coupling. We also derived [15] the Fermi liquid result:

$$E_{\mathrm{GS}}(\mathcal{R}, J'_K) = e_0 + e_1\mathcal{R} - \frac{\pi v}{24\mathcal{R}} + \left(e_2 + \frac{\pi^2 v\xi_K}{48}\right)\frac{1}{\mathcal{R}^2}, \quad (12)$$

(where e_0, e_1 and e_2 are some constants) and used it to extract $\xi_K(J'_K)$ from DMRG results for the ground state energy. Finally, we compared the magnetization versus field of our spin model to exact Bethe ansatz results [20] for the $1D$ fermion model with length $\mathcal{R}$, obtaining excellent agreement and determining ξ_K [15]. We show all our results for ξ_K versus $1/J'_K$ in Fig. 5(b) observing a good agreement between the different estimates and, at small J'_K, with the weak coupling RG prediction.

We note that, if we take region A to be the impurity spin only, then the resulting entanglement entropy can be expressed in terms of the ground state impurity magnetization, m_i: $s_{imp} = -\sum_{\pm}(1/2 \pm m_{imp})\ln(1/2 \pm m_{imp})$. (Here m_i is only non-zero when the total number of spins is odd, in which case the ground state is a spin doublet.) m_{imp} was analyzed in [8] for the $1D$ fermion Kondo model and we find similar behavior for the spin chain model. This immediately determines the behavior of s_{imp} [15].

In conclusion we have defined an impurity entanglement entropy, calculated it using DMRG and CFT methods and shown it to be a universal scaling function. We have also verified that the spin chain with a weak boundary link is indeed a realization of the Kondo model and determined accurately the corresponding Kondo length scale (ξ_K) as a function of this boundary coupling.



FIG. 5: (a) DMRG results for $S_{\mathrm{imp}}(r)\mathcal{R}/\xi_K$ for the spin chain at J_2^c for various values of J'_K in the Fermi liquid regime. The black curve is the FLT prediction, Eq. (11). (b) The various estimates for ξ_K compared with the RG result $\xi_K \propto \exp(1.376/J'_K)/\sqrt{J'_K}$.

Acknowledgements — We are grateful to J. Cardy for interesting discussions. This research was supported by NSERC (all authors), the CIAR (I.A.) CFI (E.S.) and SHARCNET (E.S.). Numerical simulations have been performed on the WestGrid network and the SCHARCNET facility at McMaster University.

---

[1] C.H. Bennett, D.P. DiVincenzo, Nature **404**, 247 (2000).
[2] See for example, G. Vidal, Phys. Rev. Lett. **93**, 040502 (2004); F. Verstraete and J.I. Cirac, cond-mat/0407066.
[3] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett. **90**, 227902 (2003); A. Osterloh, L. Amico, G. Falci, and R. Fazio, Nature (London) **4126**, 608 (2002).
[4] P. Calabrese and J. Cardy, J. Stat. Mech. 06002 (2004).
[5] G. C. Levine, Phys. Rev. Lett. 93, 266402 (2004); J. Zhao, I. Peschel and X. Wang, Phys. Rev. B **73**, 024417 (2006); H.-Q. Zhou, T. Barthel, J. Fjaerestad and U. Schollwöck, cond-mat/0511732: S.Y. Cho and R.H McKenzie, Phys. Rev. A **73**, 012109 (2006).
[6] See for example A.C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge Univ. Press. 2000).
[7] S. Eggert and I. Affleck, Phys. Rev. B **46**, 10866 (1992). S, Eggert, *ibid.*, **54**, R15590 (1996).
[8] E.S. Sørensen and I. Affleck, Phys. Rev. B **53**, 9153 (1996).
[9] V. Barzykin and I. Affleck, Phys. Rev. B **57**, 432 (1998).
[10] I. Affleck and P. Simon, Phys. Rev. Lett. **86**, 2854 (2001); E.S. Sørensen and I. Affleck, Phys. Rev. Lett. **94**, 086601 (2005); J. Gan, J. Phys. Condens. Matter **6**, 4547 (1994).
[11] M. Christandl, N. Datta, A. Ekert and A. J. Landahl, Phys. Rev. Lett. **92**, 187902 (2004), J. Zhang *et al.*, Phys. Rev. A, **72**, 012331 (2005).
[12] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher Phys. Rev. A **53**, 2046 (1996).

[13] N. Laflorencie, E.S. Sørensen, M.-S. Chang and I. Affleck, Phys. Rev. Lett. **96**, 100603 (2006).

[14] C. Holzhey, F. Larsen and F. Wilczek, Nucl. Phys. B, **424**, 443 (1994).

[15] E.S. Sørensen, M.-S. Chang, N. Laflorencie and I. Affleck, in preparation.

[16] P. Nozière, J. Low Temp. Phys. **17**, 31 (1974).

[17] I. Affleck, Nucl. Phys. B **336**, 517 (1990); I. Affleck and A.W.W. Ludwig, Nucl. Phys. B **352**, 517 (1990); ibid **B360**, 641 (1991).

[18] See for example, G. Refael and J. E. Moore, Phys. Rev. Lett. **93**, 260602 (2004).

[19] J. Cardy, J. Phys. A Math. Gen. **19**, L1093 (1986); I. Affleck, D. Gepner, H.J. Schulz, and T. Ziman, J. Phys. A Math. Gen. **22**, 511 (1989).

[20] N. Andrei, Phys. Rev. Lett. **45**, 379 (1980). P.B. Wiegman, JETP Lett., **31**, 364 (1980).